

GUIDANCE ON INFECTION CONTROL IN SCHOOLS

RASHES AND	PERIOD TO BE KEPT AWAY FROM	COMMENTS
SKIN	SCHOOL	
INFECTIONS		
Chicken pox	Until all blisters have crusted over	
German Measles	5 days from onset of rash	
Hand, foot and mouth	None	Remind children of good hygiene practice
Impetigo	48 hours after beginning antibiotic treatment	If lesions are weeping keep the child at home
Measles *	4 days from the onset of rash	
Ringworm	As soon as treatment administered	Medication is required & keep feet covered
Scabies*	Until treatment has been completed	All household and other very close contacts should be treated
Scarlet fever *	24 hours after beginning antibiotic treatment	Remind children of good hygiene practice
Slapped cheek	None	Individuals are no longer infectious once rash appears

DIARRHOEA AND VOMITING	PERIOD TO BE KEPT AWAY FROM SCHOOL	COMMENTS
Diarrhoea and/or Vomiting	48 hours from the last episode	Diarrhoea is defined as 3 or more liquid or semiliquid stools within a 24-hour period in adults and older children or any change in bowel pattern in young children. Remind children of good hygiene practice.

RESPIRATORY INFECTIONS (inc. COVID-19)	PERIOD TO BE KEPT AWAY FROM SCHOOL	COMMENTS
Includes: Common cold, COVID-19, Flu and respiratory syncytial virus (RSV)	temperature and are well enough to attend the setting.	Mild symptoms such as a runny nose, sore throat, or mild cough, who are otherwise well, should attend as normal. Remind children of good hygiene practice.
	If COVID-19, child must isolate for 3 days after the day they took their test.	
Whooping Cough		Children can still return after appropriate time, even if they are still coughing

OTHER INFECTIONS	PERIOD TO BE KEPT AWAY FROM SCHOOL	COMMENTS
Conjunctivitis	None	Remind children of good hygiene practice
Glandular fever	None	Remind children of good hygiene practice
Head lice	None	Regular head checks and treatment required
Threadworms	None	Remind children of good hygiene
Tonsillitis	None	Most cases are viruses